

On the surface gravity of a Maxwell-dilaton black hole in string theory

C S Lopez-Monsalvo*

*Instituto de Ciencias Nucleares
Universidad Nacional Autónoma de México,
A. P. 70-543, México D. F. 04510, México*

F Nettel†

*Departamento de Física, Fac. de Ciencias
Universidad Nacional Autónoma de México,
A. P. 50-542, México D. F. 04510, México*

H Quevedo‡

*Instituto de Ciencias Nucleares
Universidad Nacional Autónoma de México,
A. P. 70-543, México D. F. 04510, México*

(Dated: 19th September 2012)

The thermodynamics of dirty (Maxwell-Dilaton) black holes has been extensively studied. It has served as a fertile ground to test ideas about temperature through various definitions of surface gravity. In this paper, we make an independent analysis of this black hole solution in both, Einstein and Jordan, frames. We show that the conformal transformation between these two frames is not well defined in the extremal limit. We explore a set of definitions for surface gravity and observe the different predictions they make for the near extremal configuration of this black hole. Finally, motivated by the singularity structure in the interior of the event horizon in the Jordan frame, we use a holographic argument to remove the micro-states from the disconnected region of this solution. In this manner, we obtain a temperature which agrees with the standard results in the non-extremal regime, and which has a desirable behaviour around the extremal configurations.

* cesar.slm@correo.nucleares.unam.mx

† fnettel@ciencias.unam.mx

‡ quevedo@nucleares.unam.mx

I. INTRODUCTION

What is the temperature of an extremal configuration of a black hole spacetime? The link between thermodynamics and relativity has opened up new roads in our quest to probe gravity at a more fundamental level. Over the past thirty years a lot of work has been made to understand the connection between the laws of thermodynamics and gravitational phenomena [1–3]. This is possible through the identification of the surface gravity at the horizon of a black hole spacetime with the thermodynamic temperature of such a system [1]. One can safely use this argument in the case of stationary - and even quasi-stationary - spacetimes, in agreement with the local equilibrium hypothesis of equilibrium thermodynamics. However, the evaporation process of a near-extremal configuration of a black hole seems to be a highly dynamical setting and therefore, one is no longer entitled to use the analogy with equilibrium thermodynamics [4]. Moreover, it has been argued that extremal configurations of a black hole correspond to second order phase transitions [5]. Thus, in this context, our opening question is equivalent to that of asking about the temperature of a phase transition, taking us beyond the scope of equilibrium thermodynamics.

In this paper we address the puzzle posed by Garfinkle et-al [6] concerning the temperature of the extremal configuration of a charged black hole of string theory when working in the Jordan frame. In their article, they found the Hawking temperature of the solution through Euclidean techniques. The striking feature is that the temperature does not depend on the charge of the black hole. Thus, it is not sensitive to the near extremal configuration and the traditional arguments invoking the third law of thermodynamics to prevent the formation of a naked singularity cannot be used. To go around this issue, Garfinkle et-al claimed that the flux radiation at infinity goes to zero as the black hole becomes extremal. In such case, a discontinuous jump in the temperature from a finite value to zero takes place. This rather unusual behaviour has been used to test alternative definitions of surface gravity which are more suitable to study similar situations. In particular Hayward [7], Nielsen-Visser [8] definitions are ‘well-behaved’ at the extremal point.

Leaving aside an attempt to provide an alternative definition of surface gravity for a generic horizon, we analyse the thermodynamics of the solution found by Garfinkle et-al, in both, Einstein and Jordan, frames. As we will see the conformal factor diverges at the extremal configuration, and one should not expect to find the same physical features in both frames [9]. In particular, the peculiar singularity structure of the the Jordan frame solution suggests that a holographic argument may be used to provide a suitable expression for the entropy of the connected region of this spacetime. This allows us to find a thermodynamic temperature which complies with the requirements expected from the third law of thermodynamics and with the phase transition interpretation of the extremal configuration.

This paper is organised as follows. In section II we present the black hole solution obtained by [6, 10–12] in both Einstein and Jordan frames. We take special care of the extremal configurations, noting that the conformal factor relating these two frames diverges in this case. We found that there is a note worthy misprint in the metric for the Jordan frame in [6]. This affects the extremal configuration and alters the location of the event horizon [c.f. (21)]. We observe that the extremal configuration of the Jordan frame solution cannot be the limit of a sequence of solutions. In section III we compare the results obtained from calculating the surface gravity using a comprehensive set of definitions. We argue that the standard definition does not yield a physical result since it is unaware of the dynamical nature of the near extremal configurations. Furthermore, the non-trivial singular structure of the Jordan frame solution suggests the use of a holographic argument to remove the micro-states disconnected from the region of spacetime we analyse. Finally, we compare the different predictions from the different definitions for the near extremal behaviour.

II. A CHARGED BLACK HOLE IN STRING THEORY

Let us revise the charged low energy solution found by Gibbons and Maeda [11] and independently by Garfinkle, Horowitz and Strominger [6]. For later reference, we call this solution GMGHS. We consider the solution in both, Einstein and Jordan, frames individually, thus avoiding any possible ambiguity. The corresponding actions we are dealing with are:

1. In the Einstein frame

$$I_E = \int d^4x \sqrt{-g} \left[-R + 2(\nabla\phi)^2 + e^{-2\phi}F^2 \right], \quad (1)$$

2. and in the Jordan frame

$$I_J = \int d^4x e^{-2\phi} \sqrt{-\tilde{g}} \left[-\tilde{R} - 4\left(\tilde{\nabla}\phi\right)^2 + F^2 \right]. \quad (2)$$

Here, R is the curvature scalar, ϕ is a dilaton scalar field and F is the Faraday tensor and $\tilde{g}_{ab} = e^{2\phi}g_{ab}$.

Let us note that at this level it cannot be said that the physics derived from this two actions is the same, as it is usually assumed [9]. We need to ensure that the conformal factor is well defined everywhere in order to exchange the results derived from these two different actions. We will shortly see that the conformal factor is divergent when the solution reaches the extremal configuration, thus the transformation is not well defined.

The field equations for the Einstein frame are found in [6] and are given by

$$\nabla_a (e^{-2\phi} F^{ab}) = 0, \quad (3)$$

$$\nabla^2\phi + \frac{1}{2}e^{-2\phi}F^2 = 0, \quad (4)$$

$$R_{ab} = 2\nabla_a\phi\nabla_b\phi + 2e^{-2\phi}F_{ac}F_b{}^c - \frac{1}{2}g_{ab}e^{-2\phi}F^2, \quad (5)$$

whilst those derived from the Jordan action are

$$\tilde{\nabla}_a (e^{-2\phi} F^{ab}) = 0, \quad (6)$$

$$\tilde{\nabla}^2\phi - 2\left(\tilde{\nabla}\phi\right)^2 - \frac{1}{4}F^2 = 0, \quad (7)$$

$$\tilde{R}_{ab} = -4\tilde{\nabla}_a\phi\tilde{\nabla}_b\phi + 2F_{ac}F_b{}^c - \frac{1}{2}\tilde{g}_{ab}F^2. \quad (8)$$

A. Einstein frame solution

The field equations derived from the action (1) have the static and spherically-symmetric solution given by [12]

$$ds_E^2 = - \left[1 - \frac{2M}{r} \right] dt^2 + \left[1 - \frac{2M}{r} \right]^{-1} dr^2 + r \left[r - \frac{Q^2 e^{-2\phi_0}}{M} \right] d\Omega^2, \quad (9)$$

$$e^{-2\phi_E} = e^{-2\phi_0} \left[1 - \frac{Q^2 e^{-2\phi_0}}{Mr} \right] \quad (10)$$

and

$$F_E = Q \sin \theta d\theta \wedge d\varphi. \quad (11)$$

Here M represents the mass of the black hole, Q is the magnetic charge and ϕ_0 corresponds to the asymptotic value of the dilaton field. Note that the non-angular part of the solution is completely equivalent to that of the Schwarzschild solution. However, this similarity is only apparent since the radial coordinate does not correspond to the areal radius.

To make a thermodynamic analysis of this solution, we transform the above metric to areal coordinates. To avoid confusion and introducing unnecessary notation we keep the labels t and r of the original coordinate system. The reader should remember this point whenever using any of the following results directly on the original solution (9)-(11). Thus, using the fact that any static, spherically symmetric metric has the general form [8]

$$ds_E^2 = -\sigma(r)^2 f(r) dt^2 + f(r)^{-1} dr^2 + r^2 d\Omega^2, \quad (12)$$

we see that, in the case of (9) we have

$$f(r) = \frac{1}{4} \left[1 - \frac{4M^2 e^{2\phi_0}}{Q^2 + \sqrt{Q^4 + 4e^{4\phi_0} M^2 r^2}} \right] \left[\frac{Q^4 + 4e^{4\phi_0} M^2 r^2}{e^{4\phi_0} M^2 r^2} \right] \quad (13)$$

and

$$\sigma(r) = \frac{2Mr e^{2\phi_0}}{\sqrt{Q^4 + 4e^{4\phi_0} M^2 r^2}}. \quad (14)$$

This solution has a true curvature singularity at $r = 0$ and the event horizon is located at

$$r_+ = \sqrt{e^{-2\phi_0} [4M^2 e^{2\phi_0} - 2Q^2]}. \quad (15)$$

Thus, we see that in contrast to the Reissner-Nordström case, there is no inner horizon. The ‘extremal’ configuration occurs when the horizon completely vanishes, that is when

$$Q^2 = 2e^{2\phi_0} M^2, \quad (16)$$

and the metric becomes

$$ds_E^2 = - \frac{\sqrt{M^2 + r^2} - M}{\sqrt{M^2 + r^2} + M} dt^2 + \left[\frac{r^2}{M^2 + r^2} \right] \left[\frac{\sqrt{M^2 + r^2} + M}{\sqrt{M^2 + r^2} - M} \right] dr^2 + r^2 d\Omega^2, \quad (17)$$

which describes a naked singularity.

It follows from the dilaton solution (10) that the function $e^{2\phi}$ evaluated at the horizon diverges in the extremal limit. This is a crucial point since this function is the conformal factor relating the two frames [6]. Therefore, we cannot state that the dynamics of these two frames are equivalent in the extremal case.

B. Jordan frame solution

The metric solution of the equations of motion derived from action (2) is

$$ds_J^2 = -\frac{1 - \frac{2Me^{\phi_0}}{\rho}}{1 - \frac{Q^2 e^{-\phi_0}}{M\rho}} d\tau^2 + \frac{d\rho^2}{\left(1 - \frac{2Me^{\phi_0}}{\rho}\right)\left(1 - \frac{Q^2 e^{-\phi_0}}{M\rho}\right)} + \rho^2 d\Omega^2. \quad (18)$$

This solution can also be found through a conformal transformation of the metric (9) followed by a rescaling of the non-angular coordinates given by $\tau = e^{\phi_0} t$ and $\rho = e^{\phi_0} r$, where the temporal and spatial coordinates are labelled by τ and ρ , respectively.

The solution for the electromagnetic field is given by

$$F_J = [e^{2\phi_0} Q \sin \theta] \left[1 - \frac{Q^2 e^{-\phi_0}}{M\rho}\right]^{-1} d\theta \wedge d\varphi. \quad (19)$$

while for the dilaton field it becomes a formal expression involving integrals of Heun confluent functions.

It is straightforward to see that the metric for the extremal configuration takes the form

$$ds_J^2 = -d\tau^2 + \left[1 - \frac{2Me^{\phi_0}}{\rho}\right]^{-2} d\rho^2 + \rho^2 d\Omega^2. \quad (20)$$

Notice that in the original work of Garfinkle et-al [6] there is a misprint in the exponential factor accompanying Q^2 . In their work it appears to be $e^{3\phi_0}$ where we have $e^{-\phi_0}$. This seemingly harmless change is important since the extremal limit changes drastically, as one can verify directly if one uses their expression. However, the extremal metric, equation (15) in [6], is the correct limit of (18), above.

The horizon is located at

$$\rho_+ = 2Me^{\phi_0}, \quad (21)$$

and quick inspection of the Ricci scalar reveals that, in addition to the singularity at the origin, the curvature diverges where the inner horizon would have been, i.e. at

$$\rho_- = \frac{Q^2 e^{-\phi_0}}{M}. \quad (22)$$

This separates the spacetime into two disconnected regions divided by a singular shell. Moreover, in this case ρ is a true areal coordinate and the event horizon at $\rho = \rho_+$ depends only upon the mass of the black hole. Therefore, if we make Hawking's identification of entropy with a quarter of the area of the event horizon, and take the mass and charge as the other two thermodynamic degrees of freedom, we would not obtain a consistent first law, as we will see in the next section.




Figure 1. This figure shows the curvature scalar of a sequence of neighbouring solutions to the extremal configuration of the GMGHS black hole. The red line represents the curvature of the extremal solution, whereas the blue and green are those for which $M = M_{\text{extremal}} \pm \epsilon$. These lines diverge at the value $\rho = \rho_-$, except for the extremal one. Here, we are using $Q = 1$ and $\phi_0 = 0$ for the charge and dilaton, respectively. We take $\epsilon = 0.05, 0.005$ and 0.001 from left to right.

The extremal configuration occurs when

$$Q^2 = 2e^{2\phi_0} M^2, \quad (23)$$

and has the feature of removing the shell singularity from the solution. To see this, let us explore the curvature scalar of neighbouring solutions of the extremal configuration (20). Here, there is no longer an event horizon and all that is left is a naked singularity at the origin. In figure 1 we show a sequence of plots for the curvature scalar of various neighbouring solutions of the extremal configuration. We observe that the extremal metric is not a cluster point in the space of solutions. Moreover, no perturbation of the extremal configuration reproduces the singularity structure of the Jordan frame metric, equation (18).

From the above discussion it is clear that we are dealing with two solutions which fail to be conformally related in the extremal case. In the coming sections, we make a thermodynamic analysis for each solution separately. In particular, we will see that the two solutions represent very different thermodynamic settings.

III. SURFACE GRAVITY OF THE GMGHS BLACK HOLE

In the traditional physics jargon, surface gravity has become a synonym of temperature. Indeed, in stationary configurations, the surface gravity is an invariant quantity unambiguously defined once a normalisation convention is set. However, in a more dynamical setting a suitable definition of surface gravity becomes a more subtle matter; just as in non-equilibrium thermodynamics we can no longer define temperature without ambiguity. In this section we recall a set of definitions of surface

gravity obtained by various authors and compare their near extremal behaviour in the GMGHS solution. We proceed as in the previous section and make a separate thermodynamic analysis in each frame. Here, we limit ourselves to explore how different definitions of surface gravity match the empirical behaviour one would expect if the laws of black hole mechanics could be extended into the non-equilibrium regime. In particular, the peculiar nature of the solution in hand provides us with a useful probe to test various ideas on the matter.

The definitions of surface gravity that we use are all valid for spherical symmetry. Our selection is by no means exhaustive but we have chosen a few representatives to illustrate the different physical predictions arising from a seemingly harmless ambiguity, which we explain below.

The standard procedure to find the surface gravity of a stationary spacetime is through its timelike Killing vector field, ξ^a . At the horizon, this vector field becomes null and the surface gravity is defined as the proportionality factor measuring its departure from a geodesic curve,

$$\xi^b \nabla_b \xi_a = \kappa \xi^a. \quad (24)$$

It is conventional to adopt a unit normalisation at infinity in order to remove the ambiguity arising from rescaling of the null Killing vector. Note that this is valid only in the case of asymptotically flat spacetimes. In such case, the surface gravity is simply given by [13]

$$\kappa_1 = \left[-\frac{1}{2} \nabla_a \xi_b \nabla^a \xi^b \right]^{1/2}. \quad (25)$$

In the case of spherical symmetry there is a number of ways in which one can calculate the surface gravity. If the spacetime is also static, all of these should be equivalent. Indeed this is true for Schwarzschild and Reissner-Nordström black holes but not in general, as we show below.

Let us begin with one of the most widely used definitions of surface gravity for a static and spherically symmetric spacetime [13]

$$\kappa_2 = \lim_{r \rightarrow r_h} \left[\frac{1}{2\sqrt{-g_{tt}g_{rr}}} \frac{\partial g_{tt}}{\partial r} \right]. \quad (26)$$

In the case of a dilaton black hole (dirty black hole), Visser has shown that an equivalent way of calculating the surface gravity is given by [14]

$$\kappa_3 = \frac{1}{2r_h} e^{-\phi(r_h)} [1 - b'(r_h)], \quad (27)$$

where the function $b(r)$ is obtained by expressing the time and radial components of the metric in the form

$$g_{tt} = e^{-2\phi(r)} \left[1 - \frac{b(r)}{r} \right] \quad \text{and} \quad g_{rr} = \left[1 - \frac{b(r)}{r} \right]^{-1}, \quad (28)$$

and where the prime denotes differentiation with respect to r .

There is a more general definition introduced, by Hayward [7], valid also in the dynamical, but still spherically symmetric, scenario. It is obtained by considering the geometry of the two-surface formed by the time and radial coordinates of a spherically symmetric spacetime, and it is given by

$$\kappa_4 = \frac{1}{2} \star_2 d \star_2 dr, \quad (29)$$

where $\star_2$ represents the Hodge dual operator of the two dimensional space normal to the spheres of symmetry.

Alternatively, one can use Euclidean techniques to find the surface gravity. In this case, one starts by transforming the metric into Kruskal-Szekeres type coordinates and identifying the period of the Euclidean section in the last step of the coordinate transformation. This elegant approach, however, is not free of ambiguities either [15]. Let us write the final coordinate relation as

$$i\bar{t} = F(r) \sin\left(\frac{1}{2r_g}t\right) \quad \text{and} \quad \bar{x} = F(r) \cos\left(\frac{1}{2r_g}t\right) \quad (30)$$

where r_g denotes the gravitational radius and the Euclidean section radial functions are

$$F_E(r) = (\cosh \alpha_E + \sinh \alpha_E) \left[Q^2 e^{-2\phi_0} + \sqrt{4M^2 r^2 + Q^4 e^{-4\phi_0}} - 4M^2 \right]^{M \frac{r_+ - r_-}{r_+^2}}, \quad (31)$$

$$F_J(\rho) = (\cosh \alpha_J + \sinh \alpha_J) \sqrt{\rho - 2M e^{\phi_0}}, \quad (32)$$

with

$$\alpha_E = \frac{e^{\phi_0}}{4M} \sqrt{\frac{4M^2 r^2 + Q^4 e^{-4\phi_0}}{4M^2 e^{2\phi_0} - 2Q^2}} \quad \text{and} \quad \alpha_J = \frac{1}{4} \frac{\rho}{M e^{\phi_0}} \quad (33)$$

for the Einstein and Jordan frames, respectively. We can read out the period β of the Euclidean section directly from (30). This will be proportional to the temperature of the black hole and we can define the Euclidean surface gravity as

$$\kappa_5 = \frac{2\pi}{\beta} = \frac{1}{2r_g}, \quad (34)$$

which is equivalent to that of the Schwarzschild and Reissner-Nordström cases. This, however, is an artifact of the freedom we have in choosing the argument of the exponential functions for the Kruskal-Szekeres coordinates. In this sense, such a choice is equivalent to the normalisation at infinity in the standard definition (25).

For comparison, let us consider also the surface gravity of the Reissner-Nordström black hole [16, 17]

$$\kappa_6 = \frac{1}{2} \frac{r_+ - r_-}{r_+^2}. \quad (35)$$

Interestingly, for the GMGHS solution in the Einstein frame the value of κ_6 coincides with that of the Euclidean surface gravity, κ_5 , while in the Jordan frame it is equal to the dynamic surface gravity κ_4 .

One can also use the first law of thermodynamics to find temperature, and hence the surface gravity. To do this, we apply Hawking's expression for the entropy and use it as the thermodynamic fundamental relation, whose degrees of freedom are M and Q . Thus, the 'thermodynamic' surface gravity is given by

$$\kappa_7 = 2\pi \left[\frac{\partial S}{\partial M} \right]^{-1}. \quad (36)$$

It is important to note that this last expression is well defined only when a true thermodynamic fundamental relation is provided. That is, when the thermodynamic potential depends *at least* on two degrees of freedom. In this case, if we take the mass as our thermodynamic potential, we must have $M = M(S, Q)$ or, equivalently, $S = S(M, Q)$. In the Einstein frame the entropy is given by

$$S_E = \frac{1}{4}A_E = \pi r_+^2 = [4M^2 - 2Q^2 e^{-2\phi_0}] \pi, \quad (37)$$

where we have used the areal coordinates of the metric (12). However, when the same argument is applied in the Jordan frame we obtain

$$S_J = \frac{1}{4}A_J = \pi \rho_+^2 = 4\pi M^2 e^{2\phi_0}. \quad (38)$$

This is not a good fundamental relation and we cannot write down a Gibbs one-form for the first law of thermodynamics in the entropy representation. Nevertheless, the thermodynamic surface gravity can be easily calculated, although we cannot expect it would be of any physical relevance. Also, note that the fundamental relation, equation (37), is a separable function, and thus the temperature does not depend on Q .

There is one final point we need to attend. As we have shown in the previous section, the solution in the Jordan frame has a non-trivial singularity structure, dividing the GMGHS spacetime into two disconnected regions. All the volume enclosed by the singular shell at $\rho = \rho_-$ cannot be connected in any physical way to the exterior of the shell. Therefore, one might be over-counting the internal micro-states in the expression for the entropy (38). To make an appropriate use of the holographic principle we need to remove those microscopic degrees of freedom which cannot have an imprint in the event horizon. The simplest way of doing this, is by removing the region enclosed by the surface of radius ρ_- centred at the origin, and define an effective area enclosing the volume of the region $\rho_- < \rho < \rho_+$. Since the minimal area enclosing a given volume is a sphere, we have

$$\tilde{A}_J = 4\pi r_{\text{eff}}^2 = 4\pi (\rho_+^3 - \rho_-^3)^{\frac{2}{3}}, \quad (39)$$

where the quantity r_{eff} is the effective radius of the sphere and the associated entropy for the Jordan frame solution becomes

$$\tilde{S}_J = \pi e^{2\phi_0} \frac{[8M^6 - Q^6 e^{6\phi_0}]^{\frac{2}{3}}}{M^2}, \quad (40)$$

providing us with a true fundamental relation $\tilde{S}_J = \tilde{S}_J(M, Q)$. In this case, the thermodynamic surface gravity, equation (36), can be written in the compact form

$$\tilde{\kappa}_8 = M \frac{[\rho_+^3 - \rho_-^3]^{\frac{1}{3}}}{\rho_+^3 + \rho_-^3}. \quad (41)$$

The outcome of the different definitions presented above is summarised in table I. It is worth noting that κ_1 , κ_2 , κ_4 and κ_7 should all agree. It is interesting, however, that we do find unexpected disagreements, as discussed by Hayward et al. in [7]. In the next section we will explore the thermodynamic implications of these results.

	Einstein frame	Jordan frame
κ_1	$[4M]^{-1}$	$[4Me^{\phi_0}]^{-1}$
κ_2	$[4M]^{-1}$	$[4Me^{\phi_0}]^{-1}$
κ_3	$[4M]^{-1}$	$[4Me^{\phi_0}]^{-1}$
κ_4	$\frac{1}{8} [4M^2 e^{2\phi_0} - Q^2] [M^2 e^{\phi_0} \sqrt{4M^2 e^{2\phi_0} - 2Q^2}]^{-1}$	$\frac{1}{8} [2M^2 e^{2\phi_0} - Q^2] [M^3 e^{3\phi_0}]^{-1}$
κ_5	$\frac{1}{2} e^{\phi_0} [\sqrt{4M^2 e^{2\phi_0} - 2Q^2}]^{-1}$	$[4Me^{\phi_0}]^{-1}$
κ_6	$\frac{1}{2} e^{\phi_0} [\sqrt{4M^2 e^{2\phi_0} - 2Q^2}]^{-1}$	$\frac{1}{8} [2M^2 e^{2\phi_0} - Q^2] [M^3 e^{3\phi_0}]^{-1}$
κ_7	$[4M]^{-1}$	$[4Me^{2\phi_0}]^{-1}$
κ_8	—	$M^3 e^{2\phi_0} [8M^6 e^{6\phi_0} - Q^6]^{\frac{1}{3}} [8M^6 e^{6\phi_0} + Q^6]^{-1}$

Table I. This table shows the different results obtained after calculating the surface gravity in each frame. We observe that: (a) For Einstein frame κ_1 is in complete agreement with that obtained through Gibbs thermodynamics, κ_7 , albeit being completely featureless at the extremal configuration (see figure 2). The dynamical definition, κ_4 , shows a more interesting behaviour in the near extremal regime, where the stationary regime starts to fail. (b) For the Jordan frame, the agreement occurs between κ_4 and κ_5 , whereas κ_1 and κ_7 differ in the asymptotic value of the dilaton field. There is an extra definition, κ_8 , which corresponds to the effective Gibbs surface gravity for this solution.

IV. THERMODYNAMICS OF THE NEAR EXTREMAL CONFIGURATION

The results from the previous section show a minor disagreement between some of the definitions of surface gravity. As it was pointed out by Hayward [18], when such a situation occurs, we should make use of additional physical input to make a clear disambiguation. In figure 2 we show the temperature associated with each one of the surface gravities in table I.

In the Einstein frame, the standard definition of surface gravity, κ_1 , and that obtained through the fundamental relation (37), κ_7 , are the same. However, they are not sensitive to the extremal catastrophe which takes place, i.e. the transition from a black hole to a naked singularity. In both cases, the temperature continues to raise and diverges when all the mass has evaporated, similar to the Schwarzschild case. On the other hand, Hayward's dynamical surface gravity, κ_4 , and the RN analogous expression, κ_6 , do show some critical behaviour near the extremal point. The divergence in the extremal configuration should not be taken at face value. It is simply an indication that we are no longer in a regime where stationarity holds or, equivalently, where the local equilibrium hypothesis begins to fail. Here, small changes in the mass produce drastic changes in the thermodynamic variables. Far from the extremal regime, all definitions asymptotically agree.

At first sight, the Jordan frame solution keeps some resemblance to its counterpart but, given its different mathematical structure, there are very obvious differences. The first one, is that the standard and Gibbsian definitions of surface gravity do not exactly match. In fact, it would be very surprising if they did since the temperature associated with κ_7 is not well defined, i.e. the fundamental relation has only one thermodynamic degree of freedom [see equation (38)]. However, it comes as a surprise that the dynamical surface gravity κ_4 is actually identical to the one obtained through its Reissner-Nordström analogue. Similarly to the other frame, Hayward's definition, κ_4 , is aware of the dynamical nature surrounding the extremal point. Moreover, since it is identically zero at the extremal value, one can invoke the third law of thermodynamics to prevent the transition to a naked singularity after the event horizon disappears.

Finally, note that the temperature associated with κ_8 also goes to zero in the extremal limit, but with a vertical slope. Such behaviour describes better the thermodynamic situation in the vicinity of a phase transition. Similar to the Einstein frame, we see that small variations in the mass around the extremal configuration have a large impact on the temperature. This, again, indicates the failure of the local equilibrium hypothesis at the extremal point. Moreover, in this frame one could set up a criterion to quantify how close to the extremal configuration one can go before non-equilibrium effects dominate. For instance, the location of the maxima suggest that one can safely consider configurations with

$$M^2 > \frac{3}{2} Q^2 e^{-2\phi_0} \quad (42)$$

and

$$M^2 > \frac{1}{4} \left[32 + 8\sqrt{13} \right]^{1/3} Q^2 e^{-2\phi_0}, \quad (43)$$

for both κ_4 and κ_8 , respectively.

V. CLOSING REMARKS

In this paper we have analysed the low energy solution found independently by Gibbons and Maeda and Garfinkle, Horowitz and Strominger. We studied the conformal transformation between the Einstein and Jordan frames and showed that the conformal factor is not well defined in the extremal limit and, therefore, the corresponding dynamics are not equivalent in this case. This was a crucial point in our discussion since we were interested in the near extremal behaviour of the solution. Moreover, by inspecting the curvature scalar of the Jordan frame solution, we observed that the extremal metric cannot be a cluster point in the space of solutions to the Einstein field equations.

Motivated by a recent study of the thermodynamic geometry of this solution in the vicinity of the extremal configuration [19], we explored a comprehensive - although not exhaustive - set of definitions for the surface gravity. In our analysis, all definitions, but κ_4 , assume that the black hole transits quasi-statically through equilibrium points. This is not a valid assumption near the extremal configurations, as one can deduce from the temperature behaviour near this point. The ‘dynamical’ surface gravity, κ_4 , was constructed in order to avoid this extrapolation and to have a local definition of temperature. Moreover, it vanishes at the extremal configuration, in accordance with the third law of thermodynamics. Furthermore, inspired by the holographic principle and the singular structure of the Jordan frame solution, we tailored a definition for the thermodynamic surface gravity, κ_8 from an effective entropy, whose associated temperature also goes to zero in the extremal limit at an infinite rate. Such behaviour describes better the thermodynamic situation in the vicinity of a phase transition. The analysis is summarised in table I.

In sum, the equivalence between temperature and surface gravity is a delicate issue. There is no clear argument to prefer a particular definition other than our own physical intuition. We believe that an alternative solution to study these systems lies in the realm of a non-equilibrium thermodynamic theory for spacetime [20] or, ultimately, a quantum theory of gravity that provides us with a true counting of the gravitational micro-states [21].




Figure 2. Temperatures associated with the surface gravity definitions presented in table I.

ACKNOWLEDGEMENTS

FN receives support from DGAPA-UNAM (postdoctoral fellowship). CSLM would like to thank the Wigner Institute for their kind hospitality during the writing of this manuscript. This work was supported by CONACYT project No. 166391 and DGAPA-UNAM No. IN106110.

-
- [1] G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15**, 2752 (1977)
 - [2] J. M. Bardeen, B. Carter, and S. W. Hawking, Commun. math. Phys. **31**, 161 (1973)
 - [3] J. D. Bekenstein, Phys. Rev. D **7**, 2333 (Apr 1973)
 - [4] S. A. Hayward, Classical and Quantum Gravity **15**, 3147 (1998)
 - [5] R. G. Cai and Y. S. Myung, Nucl. Phys. B **495**, 339 (1997)
 - [6] D. Garfinkle, G. T. Horowitz, and A. Strominger, Phys. Rev. D **43**, 3140 (1991)
 - [7] S. A. Hayward, R. D. Criscienzo, M. Nadalini, L. Vanzo, and S. Zerbini, Class. Quantum Grav. **26**, 062001 (2009)
 - [8] M. Pielan, G. Kunstatter, and A. B. Nielsen, Phys. Rev. D **84**, 104008 (2011)
 - [9] A. Macias and A. Garcia, General Relativity and Gravitation **33**, 889 (2001), ISSN 0001-7701, 10.1023/A:1010212025682, <http://dx.doi.org/10.1023/A:1010212025682>
 - [10] G. W. Gibbons, Nucl. Phys. B **207**, 337 (1982)
 - [11] G. W. Gibbons and K. ichi Maeda, Nucl. Phys. B **298**, 741 (1988)
 - [12] D. Garfinkle, G. T. Horowitz, and A. Strominger, Phys. Rev. D **45**, 3888 (1992)
 - [13] R. M. Wald, *General Relativity* (Chicago University Press, 1984)
 - [14] M. Visser, Phys. Rev. D **46**, 2445 (1992)
 - [15] A. B. Nielsen and J. H. Yoon, Class. Quantum Grav. **25**, 085010 (2008)
 - [16] F. D. Felice and C. J. S. Clarke, *Relativity on curved manifolds* (Cambridge University Press, 1990)
 - [17] S. W. Hawking and G. F. R. Ellis, *The large scale structure of space-time* (Cambridge University Press, 1973)
 - [18] S. Mukohyama and S. A. Hayward, Classical and Quantum Gravity **17**, 2153 (May 2000), arXiv:gr-qc/9905085
 - [19] C. S. Lopez-Monsalvo, F. Nettel, and A. Sánchez, Braz. J. Phys. **42**, 1 (2012)
 - [20] C. S. Lopez-Monsalvo, “Covariant thermodynamics & relativity,” (March 2011), <http://eprints.soton.ac.uk/197299/>
 - [21] C. Rovelli, Class. Quantum Grav. **10**, 1549 (1993)